

Matchings in Higher-Order Gabriel Graphs*

Ahmad Biniaz[†]

Anil Maheshwari[†]

Michiel Smid[†]

May 16, 2015

Abstract

Given a set P of n points in the plane, the order- k Gabriel graph on P , denoted by k - GG , has an edge between two points p and q if and only if the closed disk with diameter pq contains at most k points of P , excluding p and q . We study matching problems in k - GG graphs. We show that a Euclidean bottleneck matching of P is contained in 10 - GG , but 8 - GG may not have any Euclidean bottleneck matching. In addition we show that 0 - GG has a matching of size at least $\frac{n-1}{4}$ and this bound is tight. We also prove that 1 - GG has a matching of size at least $\frac{2(n-1)}{5}$ and 2 - GG has a perfect matching. Finally we consider the problem of blocking the edges of k - GG .

1 Introduction

Let P be a set of n points in the plane. For any two points $p, q \in P$, let $D[p, q]$ denote the closed disk which has the line segment \overline{pq} as diameter. Let $|pq|$ be the Euclidean distance between p and q . The *Gabriel graph* on P , denoted by $GG(P)$, is defined to have an edge between two points p and q if $D[p, q]$ is empty of points in $P \setminus \{p, q\}$. Let $C(p, q)$ denote the circle which has \overline{pq} as diameter. Note that if there is a point of $P \setminus \{p, q\}$ on $C(p, q)$, then $(p, q) \notin GG(P)$. That is, (p, q) is an edge of $GG(P)$ if and only if

$$|pq|^2 < |pr|^2 + |rq|^2 \quad \forall r \in P, \quad r \neq p, q.$$

Gabriel graphs were introduced by Gabriel and Sokal [11] and can be computed in $O(n \log n)$ time [13]. Every Gabriel graph has at most $3n - 8$ edges, for $n \geq 5$, and this bound is tight [13].

A *matching* in a graph G is a set of edges without common vertices. A *perfect matching* is a matching which matches all the vertices of G . In the case that G is an edge-weighted graph, a *bottleneck matching* is defined to be a perfect matching in G in which the weight of the maximum-weight edge is minimized. For a perfect matching M , we denote the *bottleneck* of M , i.e., the length of the longest edge in M , by $\lambda(M)$. For a point set P , a *Euclidean bottleneck matching* is a perfect matching which minimizes the length of the longest edge.

In this paper we consider perfect matching and bottleneck matching admissibility of higher order Gabriel Graphs. The *order- k Gabriel graph* on P , denoted by k - GG , is the geometric graph which has an edge between two points p and q iff $D[p, q]$ contains at most k points of $P \setminus \{p, q\}$. The standard Gabriel graph, $GG(P)$, corresponds to 0 - GG . It is obvious that 0 - GG is plane, but k - GG may not be plane for $k \geq 1$. Su and Chang [14] showed that k - GG can be constructed in $O(k^2 n \log n)$ time and contains $O(k(n - k))$ edges. In [7], the authors proved that k - GG is $(k + 1)$ -connected.

*Research supported by NSERC.

[†]School of Computer Science, Carleton University, Ottawa, Canada.

1.1 Previous Work

For any two points p and q in P , the *lune* of p and q , denoted by $L(p, q)$, is defined as the intersection of the open disks of radius $|pq|$ centred at p and q . The *order- k Relative Neighborhood Graph* on P , denoted by k -*RNG*, is the geometric graph which has an edge (p, q) iff $L(p, q)$ contains at most k points of P . The *order- k Delaunay Graph* on P , denoted by k -*DG*, is the geometric graph which has an edge (p, q) iff there exists a circle through p and q which contains at most k points of P in its interior. It is obvious that

$$k\text{-RNG} \subseteq k\text{-GG} \subseteq k\text{-DG}.$$

The problem of determining whether a geometric graph has a (bottleneck) perfect matching is quite of interest. Dillencourt showed that the Delaunay triangulation (0 -*DG*) admits a perfect matching [10]. Chang et al. [9] proved that a Euclidean bottleneck perfect matching of P is contained in 16 -*RNG*.¹ This implies that 16 -*GG* and 16 -*DG* contain a (bottleneck) perfect matching of P . In [1] the authors showed that 15 -*GG* is Hamiltonian. Recently, Kaiser et al. [12] improved the bound by showing that 10 -*GG* is Hamiltonian. This implies that 10 -*GG* has a perfect matching.

Given a geometric graph $G(P)$ on a set P of n points, we say that a set K of points *blocks* $G(P)$ if in $G(P \cup K)$ there is no edge connecting two points in P , in other words, P is an independent set in $G(P \cup K)$. Aichholzer et al. [2] considered the problem of blocking the Delaunay triangulation (i.e. 0 -*DG*) for P in general position. They show that $\frac{3n}{2}$ points are sufficient to block $DT(P)$ and at least $n - 1$ points are necessary. To block a Gabriel graph, $n - 1$ points are sufficient, and $\frac{3}{4}n - o(n)$ points are sometimes necessary [3].

In a companion paper [6], we considered the matching and blocking problems in triangular-distance Delaunay (*TD-Delaunay*) graphs. The *order- k TD-Delaunay graph*, denoted by k -*TD*, on a point set P is the graph whose convex distance function is defined by a fixed-oriented equilateral triangle. Then, (p, q) is an edge in k -*TD* if there exists an equilateral triangle which has p and q on its boundary and contains at most k points of $P \setminus \{p, q\}$. We showed that 6 -*TD* contains a bottleneck perfect matching and 5 -*TD* may not have any. As for maximum matching, we proved that 1 -*TD* has a matching of size at least $\frac{2(n-1)}{5}$ and 2 -*TD* has a perfect matching (when n is even). We also showed that $\lceil \frac{n-1}{2} \rceil$ points are necessary and $n - 1$ points are sufficient to block 0 -*TD*. In [4] it is shown that 0 -*TD* has a matching of size $\lceil \frac{n-1}{3} \rceil$.

1.2 Our Results

In this paper we consider the following three problems: (a) for which values of k does every k -*GG* have a Euclidean bottleneck matching of P ? (b) for a given value k , what is the size of a maximum matching in k -*GG*? (c) how many points are sufficient/necessary to block a k -*GG*? In Section 2 we review and prove some graph-theoretic notions. In Section 3 we consider the problem (a) and prove that a Euclidean bottleneck matching of P is contained in 10 -*GG*. In addition, we show that for some point sets, 8 -*GG* does not have any Euclidean bottleneck matching. In Section 4 we consider the problem (b) and give some lower bounds on the size of a maximum matching in k -*GG*. We prove that 0 -*GG* has a matching of size at least $\frac{n-1}{4}$, and this bound is tight. In addition we prove that 1 -*GG* has a matching of size at least $\frac{2(n-1)}{5}$ and 2 -*GG* has a perfect matching. In Section 5 we consider the problem (c). We show that at least $\lceil \frac{n-1}{3} \rceil$ points are necessary to block a Gabriel graph and this bound is tight. We also show that at least $\lceil \frac{(k+1)(n-1)}{3} \rceil$ points are necessary and $(k+1)(n-1)$ points are sufficient to block a k -*GG*. The open problems and concluding remarks are presented in Section 6.

¹They defined k -*RNG* in such a way that $L(p, q)$ contains at most $k - 1$ points of P .

2 Preliminaries

Let G be an edge-weighted graph with vertex set V and weight function $w : E \rightarrow \mathbb{R}^+$. Let T be a minimum spanning tree of G , and let $w(T)$ be the total weight of T .

Lemma 1. *Let $\delta(e)$ be a cycle in G that contains an edge $e \in T$. Let δ' be the set of edges in $\delta(e)$ which do not belong to T and let e'_{max} be the largest edge in δ' . Then, $w(e) \leq w(e'_{max})$.*

Proof. Let $e = (u, v)$ and let T_u and T_v be the two trees obtained by removing e from T . Let $e' = (x, y)$ be an edge in δ' such that one of x and y belongs to T_u and the other one belongs to T_v . By definition of e'_{max} , we have $w(e') \leq w(e'_{max})$. Let $T' = T_u \cup T_v \cup \{(x, y)\}$. Clearly, T' is a spanning tree of G . If $w(e') < w(e)$ then $w(T') < w(T)$; contradicting the minimality of T . Thus, $w(e) \leq w(e')$, which completes the proof of the lemma. \square

For a graph $G = (V, E)$ and $S \subseteq V$, let $G - S$ be the subgraph obtained from G by removing all vertices in S , and let $o(G - S)$ be the number of odd components in $G - S$, i.e., connected components with an odd number of vertices. The following theorem by Tutte [15] gives a characterization of the graphs which have perfect matching:

Theorem 1 (Tutte [15]). *G has a perfect matching if and only if $o(G - S) \leq |S|$ for all $S \subseteq V$.*

Berge [5] extended Tutte's theorem to a formula (known as the Tutte-Berge formula) for the maximum size of a matching in a graph. In a graph G , the *deficiency*, $\text{def}_G(S)$, is $o(G - S) - |S|$. Let $\text{def}(G) = \max_{S \subseteq V} \text{def}_G(S)$.

Theorem 2 (Tutte-Berge formula; Berge [5]). *The size of a maximum matching in G is*

$$\frac{1}{2}(n - \text{def}(G)).$$

For an edge-weighted graph G we define the *weight sequence* of G , $\text{WS}(G)$, as the sequence containing the weights of the edges of G in non-increasing order. A graph G_1 is said to be less than a graph G_2 if $\text{WS}(G_1)$ is lexicographically smaller than $\text{WS}(G_2)$.

3 Euclidean Bottleneck Matching

Given a point set P , in this section we prove that 10- GG contains a Euclidean bottleneck matching of P . We also present a configuration of a point set P such that 8- GG does not contain any Euclidean bottleneck matching of P . We use a similar argument as in [1, 8]. First consider the following lemma of [1]:

Lemma 2 (Abellanas et al. [1]). *Let $0 < \theta \leq \pi/5$. Let $C(A, \theta, L, R)$ be a cone with apex A , bounding rays L and R emanating from A and angle θ computed clockwise from L to R . Consider two points $x, y \in C(A, \theta, L, R)$ and a constant $r > 0$. If $|xA| > 2r$ and $|yA| > 2r$, then $|xy| < 2r$ or $|xy| < \max\{|xA| - r, |yA| - r\}$.*

Theorem 3. *For every point set P , 10- GG contains a Euclidean bottleneck matching of P .*

Proof. Let \mathcal{M} be the set of all perfect matchings on the points of P . Define a total order on the elements of \mathcal{M} by their weight sequence. If two elements have exactly the same weight sequence, break ties arbitrarily to get a total order. Let $M^* = \{(a_1, b_1), \dots, (a_{\frac{n}{2}}, b_{\frac{n}{2}})\}$ be a perfect matching in \mathcal{M} with minimal weight sequence. It is obvious that M^* is a Euclidean bottleneck matching for P . We will show that all edges of M^* are in 10- GG . Consider any




Figure 1: Illustration for Theorem 3.

edge $e = (a_i, b_i)$ in M^* and its corresponding disk $D[a_i, b_i]$. Suppose that $D[a_i, b_i]$ contains w points of $P \setminus \{a_i, b_i\}$. Let $U = \{u_1, u_2, \dots, u_w\}$ represent the points inside $D[a_i, b_i]$, and $U' = \{r_1, r_2, \dots, r_w\}$ represent the points where $(r_i, u_i) \in M^*$. We will show that $w \leq 10$. Let $r = |a_i b_i|/2$ be the radius of $D[a_i, b_i]$.

Claim 1: For each $r_j \in U'$, $\min\{|r_j a_i|, |r_j b_i|\} \geq 2r$. To prove this, assume that $|r_j a_i| < 2r$ and let M be the perfect matching obtained from M^* by deleting $\{(a_i, b_i), (r_j, u_j)\}$, and adding $\{(a_i, r_j), (b_i, u_j)\}$. The two new edges are smaller than the old ones. Thus, $\text{WS}(M) < \text{WS}(M^*)$ which contradicts the minimality of M^* .

Let D_1 and D_2 respectively be the open disks with radius $2r$ centered at a_i and b_i . By Claim 1, we know that no point of U' lies inside $D_1 \cup D_2$. In other words all points of U' are contained in $\overline{D_1 \cup D_2}$.

Claim 2: For each pair r_j and r_k of points in U' , $|r_j r_k| \geq \max\{|a_i b_i|, |r_j u_j|, |r_k u_k|\}$. To prove this, assume that $|r_j r_k| < \max\{|a_i b_i|, |r_j u_j|, |r_k u_k|\}$. Let M be the perfect matching obtained from M^* by deleting $\{(u_j, r_j), (u_k, r_k), (a_i, b_i)\}$ and adding $\{(a_i, u_j), (b_i, u_k), (r_j, r_k)\}$. Since $\max\{|a_i u_j|, |b_i u_k|, |r_j r_k|\} < \max\{|u_j r_j|, |u_k r_k|, |a_i b_i|\}$, $\text{WS}(M) < \text{WS}(M^*)$ which contradicts the minimality of M^* .

Let c_i be the center of $D[a_i, b_i]$. Consider a decomposition of the plane into 10 cones C_1, \dots, C_{10} of angle $\pi/5$ with apex at c_i . See Figure 1. By contradiction, we will show that each cone $C_i, 1 \leq i \leq 10$, contains at most one point of U' . Suppose that a cone C_i where $1 \leq i \leq 10$ contains two points $r_j, r_k \in U'$. It is obvious that

$$|r_j u_j| \geq |r_j c_i| - r \quad \text{and} \quad |r_k u_k| \geq |r_k c_i| - r. \quad (1)$$

Claim 3: Each cone C_i where $1 \leq i \leq 10$ and $i \neq 3, 8$ contains at most one point of U' . Suppose that C_i contain two points $r_j, r_k \in U'$. By Claim 1, r_j and r_k are contained in $\overline{D_1 \cup D_2}$. Consider the disk D_3 with radius $2r$ centered at c_i , as shown in Figure 1. Since the intersection of D_3 and $\overline{D_1 \cup D_2}$ in C_i is empty, r_j and r_k are outside D_3 , i.e., $|r_j c_i| > 2r$ and $|r_k c_i| > 2r$. By Lemma 2, $|r_j r_k| < 2r$ or $|r_j r_k| < \max\{|r_j c_i| - r, |r_k c_i| - r\}$. By inequality (1), $|r_j r_k| < \max\{|a_i b_i|, |r_j u_j|, |r_k u_k|\}$ which contradicts Claim 2.

Claim 4: Each of C_3 and C_8 contains at most one point of U' . Let S_1 and S_2 be the connected components of $D_3 \cap (\overline{D_1 \cup D_2})$ which lie inside C_3 and C_8 , respectively, as shown




Figure 2: (a) The angle $\angle bac$ is smaller than the angle $\angle abc$, and hence (b) $\angle r_k a' r_j < \angle a' r_k r_j$.

in Figure 1. Because of symmetry, we only prove the claim for C_3 . Suppose that C_3 contains two points $r_j, r_k \in U'$. For the rest of the proof, refer to Figure 2. W.l.o.g. assume that r_j is further from c_i than r_k and r_k is to the left of r_j (i.e., r_k is to the left of the line through c_i and r_j oriented from c_i to r_j). If $r_k \notin S_1$ then $|r_k c_i| > 2r$ and $|r_j c_i| > 2r$. Then, by Lemma 2 and Claim 2 we have a contradiction. Therefore, assume that $r_k \in S_1$. Let L and R be the two rays defining C_3 . Let a be the intersection of R and $C(a_i, b_i)$. Let b be the intersection of the boundaries of D_1 and D_3 which is inside C_3 . Define the point c on R such that $|bc| = 2r$ and $c \neq c_i$. See Figure 2(a). The triangle $\triangle b c c_i$ is isosceles, and hence $\angle b c c_i = \angle b c_i c < \frac{\pi}{5}$. This implies that $\angle c b c_i > \frac{3\pi}{5}$. On the other hand, in triangle $\triangle a b c_i$, $|ab| > |a c_i|$, which implies that $\angle a b c_i < \angle a c_i b < \frac{\pi}{5}$. Thus $\angle a b c > \frac{2\pi}{5}$. In addition $\angle b a c_i > \frac{3\pi}{5}$ and hence $\angle b a c < \frac{2\pi}{5}$. Therefore in the triangle $\triangle a b c$ we have

$$\angle a b c > \frac{2\pi}{5} > \angle b a c.$$

Let $C(b, c)$ be the circle with radius $2r$ having \overline{bc} as diameter, and let A be the ray emanating from b which goes through c as shown in Figure 2(b). The intersection of C_3 with $\overline{D_1} \cup \overline{D_2}$ which lies to the right of A is completely inside $C(b, c)$. Thus, if r_j is to the right of A , $|r_j r_k| < 2r = |a_i b_i|$, which contradicts Claim 2. Therefore r_j lies to the left of A . If r_j is in the interior of C_3 , rotate C_3 counter-clockwise around c_i until r_j lies on R . Since r_k is to the left of r_j , the point r_k is still in the interior of C_3 . Let a' be the intersection of the new R with $C(a_i, b_i)$. Note that S_1 and hence r_k is contained in $\triangle a b c$. In addition r_j and a' are outside $\triangle a b c$ and to the left of the line through a and c . Therefore, $\angle a' r_k r_j \geq \angle a b c > \angle b a c \geq \angle r_k a' r_j$ and hence

$$|r_j r_k| < |r_j a'| = |r_j c_i| - r \leq |r_j u_j|,$$

which contradicts Claim 2.

By Claim 3 and Claim 4 each cone C_i where $1 \leq i \leq 10$ contains at most one point of U' . Thus, $w \leq 10$, and $e = (a_i, b_i)$ is an edge of $10\text{-}GG$. \square

Now, we will show that for some point sets, $8\text{-}GG$ does not contain any Euclidean bottleneck matching. Consider Figure 3 which shows a configuration of a set P of 20 points. The closed disk $D[a, b]$ is centred at c and has diameter one, i.e., $|ab| = 1$. $D[a, b]$ contains 9 points $U = \{u_1, \dots, u_9\}$ which lie on a circle with radius $\frac{1}{2} - \epsilon$ which is centred at c . Nine points in




Figure 3: A set of 20 points such that $8\text{-}GG$ does not contain any Euclidean bottleneck matching.

$U' = \{r_1, \dots, r_9\}$ are placed on a circle with radius 1.5 which is centred at c in such a way that $|r_j u_j| = 1 + \epsilon$, $|r_j a| > 1 + \epsilon$, $|r_j b| > 1 + \epsilon$, and $|r_j r_k| > 1 + \epsilon$ for $1 \leq j, k \leq 9$ and $j \neq k$. Consider a perfect matching $M = \{(a, b)\} \cup \{(r_i, u_i) : i = 1, \dots, 9\}$ where each point $r_i \in U'$ is matched to its closest point u_i . It is obvious that $\lambda(M) = 1 + \epsilon$, and hence the bottleneck of any bottleneck perfect matching is at most $1 + \epsilon$. We will show that any Euclidean bottleneck matching of P contains (a, b) . By contradiction, let M^* be a Euclidean bottleneck matching which does not contain (a, b) . In M^* , a is matched to a point $x \in U \cup U'$. If $x \in U'$, then $|ax| > 1 + \epsilon$. If $x \in U$, w.l.o.g. assume that $x = u_1$. Thus, in M^* the point r_1 is matched to a point y where $y \neq u_1$. Since u_1 is the closest point to r_1 and $|r_1 u_1| = 1 + \epsilon$, $|r_1 y| > 1 + \epsilon$. In both cases $\lambda(M^*) > 1 + \epsilon$, which is a contradiction. Therefore, M^* contains (a, b) . Since $D[a, b]$ contains 9 points of $P \setminus \{a, b\}$, $(a, b) \notin 8\text{-}GG$. Therefore $8\text{-}GG$ does not contain any Euclidean bottleneck matching of P .

4 Maximum Matching

Let P be a set of n points in the plane. In this section we will prove that $0\text{-}GG$ has a matching of size at least $\frac{n-1}{4}$; this bound is tight. We also prove that $1\text{-}GG$ has a matching of size at least $\frac{2(n-1)}{5}$ and $2\text{-}GG$ has a perfect matching (when n is even).

First we give a lower bound on the number of components that result after removing a set

S of vertices from k - GG . Then we use Theorem 1 and Theorem 2, respectively presented by Tutte [15] and Berge [5], to prove a lower bound on the size of a maximum matching in k - GG .




Figure 4: The point set P of 16 points is partitioned into white/black disks, white/black squares, and crosses. (a) The graph $G(\mathcal{P})$, (b) The set \mathcal{T} of straight-line edges corresponding to $MST(G(\mathcal{P}))$ is in bold, and the set \mathcal{D} of their corresponding disks.

Let $\mathcal{P} = \{P_1, P_2, \dots\}$ be a partition of the points in P . For two sets P_i and P_j in \mathcal{P} define the distance $d(P_i, P_j)$ as the smallest Euclidean distance between a point in P_i and a point in P_j , i.e., $d(P_i, P_j) = \min\{|ab| : a \in P_i, b \in P_j\}$. Let $G(\mathcal{P})$ be the complete edge-weighted graph with vertex set \mathcal{P} . For each edge $e = (P_i, P_j)$ in $G(\mathcal{P})$, let $w(e) = d(P_i, P_j)$. This edge e is defined by two points a and b , where $a \in P_i$ and $b \in P_j$. Therefore, an edge $e \in G(\mathcal{P})$ corresponds to a straight line edge between two points $a, b \in P$; see Figure 4(a). Let $MST(G(\mathcal{P}))$ be a minimum spanning tree of $G(\mathcal{P})$. It is obvious that each edge e in $MST(G(\mathcal{P}))$ corresponds to a straight line edge between $a, b \in P$. Let \mathcal{T} be the set of all these straight line edges. Let \mathcal{D} be the set of disks which have the edges of \mathcal{T} as diameter, i.e., $\mathcal{D} = \{D[a, b] : (a, b) \in \mathcal{T}\}$. See Figure 4(b).

Observation 1. \mathcal{T} is a subgraph of a minimum spanning tree of P , and hence \mathcal{T} is plane.

Lemma 3. A disk $D[a, b] \in \mathcal{D}$ does not contain any point of $P \setminus \{a, b\}$.

Proof. By Observation 1, \mathcal{T} is a subgraph of a minimum spanning tree of P . It is well known that any minimum spanning tree of P is a subgraph of 0 - $GG(P)$. Thus, \mathcal{T} is a subgraph of 0 - $GG(P)$, and hence, any disk $D[a, b] \in \mathcal{D}$ —representing an edge of \mathcal{T} —does not contain any point of $P \setminus \{a, b\}$. \square

Lemma 4. For each pair D_i and D_j of disks in \mathcal{D} , D_i (resp. D_j) does not contain the center of D_j (resp. D_i).

Proof. Let (a_i, b_i) and (a_j, b_j) respectively be the edges of \mathcal{T} which correspond to D_i and D_j . Let C_i and C_j be the circles representing the boundary of D_i and D_j . W.l.o.g. assume that C_j is the bigger circle, i.e., $|a_i b_i| < |a_j b_j|$. By contradiction, suppose that C_j contains the center c_i of C_i . Let x and y denote the intersections of C_i and C_j . Let x_i (resp. x_j) be the intersection of C_i (resp. C_j) with the line through y and c_i (resp. c_j). Similarly, let y_i (resp. y_j) be the intersection of C_i (resp. C_j) with the line through x and c_i (resp. c_j).

As illustrated in Figure 5, the arcs $\widehat{x_i x}$, $\widehat{y_i y}$, $\widehat{x_j x}$, and $\widehat{y_j y}$ are the potential positions for the points a_i, b_i, a_j , and b_j , respectively. First we will show that the line segment $x_i x_j$ passes through x and $|a_i a_j| \leq |x_i x_j|$. The angles $\angle x_i x y$ and $\angle x_j x y$ are right angles, thus the line segment $x_i x_j$ goes through x . Since $\widehat{x_i x} < \pi$ (resp. $\widehat{x_j x} < \pi$), for any point $a_i \in \widehat{x_i x}$, $|a_i x| \leq |x_i x|$ (resp. $a_j \in \widehat{x_j x}$, $|a_j x| \leq |x_j x|$). Therefore,

$$|a_i a_j| \leq |a_i x| + |x a_j| \leq |x_i x| + |x x_j| = |x_i x_j|.$$




Figure 5: Illustration of Lemma 4: C_i and C_j intersect, and C_j contains the center of C_i .

Consider triangle $\Delta x_i x_j y$ which is partitioned by segment $c_i x_j$ into $t_1 = \Delta x_i x_j c_i$ and $t_2 = \Delta c_i x_j y$. Since $x_i y$ is a diameter of C_i which passes through the center c_i , the length of the segment $x_i c_i$ of t_1 is equal to the length of the segment $c_i y$ of t_2 . The segment $c_i x_j$ is shared by t_1 and t_2 . Since c_i is inside C_j and $\widehat{y x_j} = \pi$, the angle $\angle y c_i x_j > \frac{\pi}{2}$. Thus, $\angle x_i c_i x_j$ in t_1 is smaller than $\frac{\pi}{2}$ (and hence smaller than $\angle y c_i x_j$ in t_2). Therefore, the segment $x_i x_j$ of t_1 is smaller than the segment $x_j y$ of t_2 . Thus,

$$|a_i a_j| \leq |x_i x_j| < |x_j y| = |a_j b_j|.$$

By symmetry $|b_i b_j| < |a_j b_j|$. Therefore $\max\{|a_i a_j|, |b_i b_j|\} < \max\{|a_i b_i|, |a_j b_j|\}$. In addition $\delta = (a_i, a_j, b_j, b_i, a_i)$ is a cycle and at least one of (a_i, a_j) and (b_i, b_j) does not belong to \mathcal{T} . This contradicts Lemma 1 (Note that by Observation 1, \mathcal{T} is a subgraph of a minimum spanning tree of P). \square

Now we show that the intersection of every four disks in \mathcal{D} is empty. In other words, every point in the plane cannot lie in more than three disks in \mathcal{D} . In Section 4.1 we prove the following theorem, and in Section 4.2 we present the lower bounds on the size of a maximum matching in k -GG.

Theorem 4. *For every four disks $D_1, D_2, D_3, D_4 \in \mathcal{D}$, $D_1 \cap D_2 \cap D_3 \cap D_4 = \emptyset$.*

4.1 Proof of Theorem 4

The proof is by contradiction. Let D_1, D_2, D_3 , and D_4 be four disks in \mathcal{D} . Let $\mathcal{X} = D_1 \cap D_2 \cap D_3 \cap D_4$ and let x be a point in \mathcal{X} . For $i = 1, 2, 3, 4$, let c_i be the center of D_i , let C_i be the boundary of D_i , and let (a_i, b_i) be the edge in \mathcal{T} which corresponds to D_i . Denote the angle $\angle a_i x b_i$ by α_i , for $i = 1, 2, 3, 4$. Since (a_i, b_i) is a diameter of D_i and x lies in D_i , $\alpha_i \geq \frac{\pi}{2}$. First we prove the following observation.

Observation 2. *For $i, j \in \{1, 2, 3, 4\}$, where $i \neq j$, the angles α_i and α_j are either disjoint or one is completely contained in the other.*

Proof. The proof is by contradiction. Suppose that α_i and α_j are not disjoint and none of them is completely contained in the other. Thus α_i and α_j share some part and w.l.o.g. assume that b_i is in the cone which is obtained by extending the edges of α_j , and b_j is in the cone which is obtained by extending the edges of α_i . Three cases arise:

- $b_i \in \Delta x a_j b_j$. In this case b_i is inside D_j which contradicts Lemma 3.
- $b_j \in \Delta x a_i b_i$. In this case b_j is inside D_i which contradicts Lemma 3.
- $b_i \notin \Delta x a_j b_j$ and $b_j \notin \Delta x a_i b_i$. In this case (a_i, b_i) intersects (a_j, b_j) which contradicts Observation 1.

□

We call α_i a *blocked angle* if α_i is contained in an angle α_j for some $j \in \{1, 2, 3, 4\}$, where $j \neq i$. Otherwise, we call α_i a *free angle*.

Lemma 5. *At least one α_i , for $i \in \{1, 2, 3, 4\}$, is blocked.*

Proof. Suppose that all angles α_i , where $i \in \{1, 2, 3, 4\}$, are free. This implies that the α_i s are pairwise disjoint and $\alpha = \sum_{i=1}^4 \alpha_i \geq 2\pi$. If $\alpha > 2\pi$, we obtain a contradiction to the fact that the sum of the disjoint angles around x is at most 2π . If $\alpha = 2\pi$, then the four edges (a_i, b_i) where $i \in \{1, 2, 3, 4\}$, form a cycle which contradicts the fact that \mathcal{T} is a subgraph of a minimum spanning tree of P . □




Figure 6: (a) The point x should be inside the circle segment $sc(c_i, b_i)$. (b) The trap (a_i, b_i) which consists of two lenses $lens(a_i, c_i)$ and $lens(b_i, c_i)$.

By Lemma 5 at least one of the angles is blocked. Hereafter, assume that α_j is blocked by α_i where $1 \leq i, j \leq 4$ and $i \neq j$. W.l.o.g. assume that $a_i b_i$ is a vertical line segment and the point x (which belongs to \mathcal{X}) is to the left of $a_i b_i$. Thus, $a_j b_j$ and c_j are to the right of $a_i b_i$. This implies that $a_i b_i \cap D_j \neq \emptyset$. See Figure 6(a). By Lemma 4, c_i cannot be inside D_j , thus either $a_i c_i \cap D_j \neq \emptyset$ or $c_i b_i \cap D_j \neq \emptyset$, but not both. W.l.o.g. assume that $c_i b_i \cap D_j \neq \emptyset$. Let C' be the circle with radius $|c_i b_i|$ which is centered at b_i . Let d denote the intersection of C' with C_i which is to the right of $c_i b_i$. Consider the circle C'' with radius $|db_i|$ centered at d . Note that C'' goes through b_i and c_i . Let $sc(c_i, b_i)$ be the segment of the circle C'' which is between the chord $c_i b_i$ and the arc $\widehat{c_i b_i}$ as shown in Figure 6(a).

We show that x cannot be outside $\text{sc}(c_i, b_i)$. By contradiction suppose that x is outside $\text{sc}(c_i, b_i)$ (and to the left of $c_i b_i$). Let l_1 and l_2 respectively be the perpendicular bisectors of $x b_i$ and $x c_i$. Let b'_i and c'_i respectively be the intersection of l_1 and l_2 with $c_i b_i$ and let d' be the intersection point of l_1 and l_2 . Since x is outside $\text{sc}(c_i, b_i)$, the intersection point d' is to the left of (the vertical line through) d and inside triangle $\triangle b_i c_i d$. If c_j is below l_1 then $|c_j b_i| < |c_j x|$ and D_j contains b_i which contradicts Lemma 4. If c_j is above l_2 then $|c_j b_i| < |c_j x|$ and D_j contains c_i which contradicts Lemma 4. Thus, c_j is above l_1 and below l_2 , and (by the initial assumption) to the right of $c_i b_i$. That is, c_j is in triangle $\triangle b'_i c'_i d'$. Since $\triangle b'_i c'_i d' \subseteq \triangle b_i c_i d \subseteq D_i$, c_j lies inside D_i which contradicts Lemma 4. Therefore, x is contained in $\text{sc}(c_i, b_i)$.

By symmetry D_j can intersect $a_i c_i$ and/or c_j can be to the left of $a_i b_i$ as well. Therefore, if a_i blocks α_j , the point x can be in $\text{sc}(c_i, b_i)$ or any of the symmetric segments of the circles. For an edge $a_i b_i$ we denote the union of these segments by $\text{trap}(a_i, b_i)$ which is shown in Figure 6(b). For each disk D_i , let $\text{trap}(D_i) = \text{trap}(a_i, b_i)$ where (a_i, b_i) is the edge in \mathcal{T} corresponding to D_i . Therefore x is contained in $\text{trap}(D_i)$ which implies that

$$\mathcal{X} \subseteq \text{trap}(D_i).$$

Note that $\text{trap}(D_i)$ consists of two symmetric lenses $\text{lens}(a_i, c_i)$ and $\text{lens}(b_i, c_i)$, i.e., $\text{trap}(D_i) = \text{lens}(a_i, c_i) \cup \text{lens}(b_i, c_i)$.

Lemma 6. For any point $x \in \text{trap}(a_i, b_i)$, $\angle a_i x b_i \geq 150^\circ$.

Proof. See Figure 6(a). The angle $\angle b_i d c_i = 60^\circ$, which implies that $\widehat{c_i b_i} = 60^\circ$. Thus, for any point x' on the arc $\widehat{c_i b_i}$, $\angle x' c_i b_i + \angle x' b_i c_i = 30^\circ$, and hence for any point x in the segment $\text{sc}(c_i, b_i)$, $\angle x c_i b_i + \angle x b_i c_i \leq 30^\circ$. This implies that in $\triangle x b_i c_i$, $\angle b_i x c_i \geq 150^\circ$. On the other hand $\angle b_i x c_i \leq \angle b_i x a_i$, which proves the lemma. \square




Figure 7: Illustration of Lemma 7.

Lemma 7. For any two disks D_i and D_j in \mathcal{D} , $\text{trap}(D_i) \cap \text{trap}(D_j) = \emptyset$.

Proof. We prove this lemma by contradiction. Suppose $x \in \text{trap}(D_i) \cap \text{trap}(D_j)$ and w.l.o.g. assume that $x \in \text{lens}(a_i, c_i) \cap \text{lens}(a_j, c_j)$ as shown in Figure 7. Connect x to a_i , c_i , a_j , and c_j (a_i may be identified with a_j). As shown in the proof of Lemma 6, $\min\{\angle a_i x c_i, \angle a_j x c_j\} > 150^\circ$. Two configurations may arise:

- $\angle c_i x c_j \leq 60^\circ$. In this case $|c_i c_j| \leq \max\{|x c_i|, |x c_j|\}$. W.l.o.g. assume that $|x c_i| \leq |x c_j|$ which implies that $|c_i c_j| \leq |x c_j|$; see Figure 7(a). Clearly $|x c_j| < |c_j a_j|$, and hence $|c_i c_j| < |c_j a_j|$. Thus, D_j contains c_i which contradicts Lemma 4.
- $\angle c_i x c_j > 60^\circ$. In this case $\angle a_i x c_j \leq 60^\circ$ and $\angle a_j x c_i \leq 60^\circ$, hence $|a_i c_j| \leq \max\{|a_i x|, |c_j x|\}$ and $|a_j c_i| \leq \max\{|a_j x|, |c_i x|\}$. Three configurations arise:
 - $|a_i x| < |c_j x|$, in this case $|a_i c_j| < |c_j x| < |c_j a_j|$ and hence D_j contains a_i . See Figure 7(b).
 - $|a_j x| < |c_i x|$, in this case $|a_j c_i| < |c_i x| < |c_i a_i|$ and hence D_i contains a_j .
 - $|a_i x| \geq |c_j x|$ and $|a_j x| \geq |c_i x|$, in this case w.l.o.g. assume that $|a_i x| \leq |a_j x|$. Thus $|a_i c_j| \leq |a_i x| \leq |a_j x| < |a_j c_j|$ which implies that D_j contains a_i . See Figure 7(b).

All cases contradict Lemma 3. □

Recall that each blocking angle represents a trap. Thus, by Lemma 5 and Lemma 7, we have the following corollary:

Corollary 1. *Exactly one α_i , where $1 \leq i \leq 4$, is blocked.*

Recall that α_j is blocked by α_i , $a_i b_i$ is vertical line segment, c_j is to the right of $a_i b_i$, and $x \in \text{sc}(c_i, b_i)$. As a direct consequence of Corollary 1, α_i , α_k , and α_l are free angles, where $1 \leq i, j, k, l \leq 4$ and $i \neq j \neq k \neq l$. In addition, c_k and c_l are to the left of $a_i b_i$. It is obvious that

$$\mathcal{X} \subseteq \text{trap}(D_i) \cap D_k \cap D_l.$$




Figure 8: Illustration of Lemma 8.

Lemma 8. For a blocking angle α_i and free angles α_k and α_l , $\text{trap}(D_i) \cap D_k \cap D_l = \emptyset$.

Proof. Since α_i is a blocking angle and α_k, α_l are free angles, c_k and c_l are on the same side of $a_i b_i$. By contradiction, suppose that $x \in \text{trap}(D_i) \cap D_j \cap D_k$. See Figure 8. It is obvious that $\max\{|xa_i|, |xb_i|\} < |a_i b_i|$, $\max\{|xa_k|, |xb_k|\} < |a_k b_k|$, and $\max\{|xa_l|, |xb_l|\} < |a_l b_l|$. By Lemma 6, $\alpha_i \geq 150^\circ$. In addition $\alpha_k, \alpha_l \geq 90^\circ$. Thus, $\max\{\angle a_i x b_k, \angle a_k x b_l, \angle a_l x b_i\} \leq 30^\circ$. Hence, $|a_i b_k| < \max\{|xa_i|, |xb_k|\}$, $|a_k b_l| < \max\{|xa_k|, |xb_l|\}$, and $|a_l b_i| < \max\{|xa_l|, |xb_i|\}$. Therefore, $\max\{|a_i b_k|, |a_k b_l|, |a_l b_i|\} < \max\{|a_i b_i|, |a_k b_k|, |a_l b_l|\}$. In addition $\delta = (a_i, b_i, a_l, b_l, a_k, b_k, a_i)$ is a cycle and at least one of (a_i, b_k) , (a_k, b_l) and (a_l, b_i) does not belong to \mathcal{T} . This contradicts Lemma 1. \square

Thus, $\mathcal{X} = \emptyset$; which contradicts the fact that $x \in \mathcal{X}$. This completes the proof of Theorem 4.

4.2 Lower Bounds

In this section we present some lower bounds on the size of a maximum matching in 2-GG, 1-GG, and 0-GG.

Theorem 5. For a set P of an even number of points, 2-GG has a perfect matching.

Proof. First we show that by removing a set S of s points from 2-GG, at most $s+1$ components are generated. Then we show that at least one of these components must be even. Using Theorem 1, we conclude that 2-GG has a perfect matching.

Let S be a set of s vertices removed from 2-GG, and let $\mathcal{C} = \{C_1, \dots, C_{m(s)}\}$ be the resulting $m(s)$ components, where m is a function depending on s . Actually $\mathcal{C} = 2\text{-GG} - S$ and $\mathcal{P} = \{V(C_1), \dots, V(C_{m(s)})\}$ is a partition of the vertices in $P \setminus S$.

Claim 1. $m(s) \leq s+1$. Let $G(\mathcal{P})$ be the complete graph with vertex set \mathcal{P} which is constructed as described above. Let \mathcal{T} be the set of all edges between points in P corresponding to the edges of $MST(G(\mathcal{P}))$ and let \mathcal{D} be the set of disks corresponding to the edges of \mathcal{T} . It is obvious that \mathcal{T} contains $m(s) - 1$ edges and hence $|\mathcal{D}| = m(s) - 1$. Let $F = \{(p, D) : p \in S, D \in \mathcal{D}, p \in D\}$ be the set of all (point, disk) pairs where $p \in S$, $D \in \mathcal{D}$, and p is inside D . By Theorem 4 each point in S can be inside at most three disks in \mathcal{D} . Thus, $|F| \leq 3 \cdot |S|$. Now we show that each disk in \mathcal{D} contains at least three points of S in its interior. Consider any disk $D \in \mathcal{D}$ and let $e = (a, b)$ be the edge of \mathcal{T} corresponding to D . By Lemma 3, D does not contain any point of $P \setminus S$. Therefore, D contains at least three points of S , because otherwise (a, b) is an edge in 2-GG which contradicts the fact that a and b belong to different components in \mathcal{C} . Thus, each disk in \mathcal{D} has at least three points of S . That is, $3 \cdot |\mathcal{D}| \leq |F|$. Therefore, $3(m(s) - 1) \leq |F| \leq 3s$, and hence $m(s) \leq s+1$.

Claim 2: $o(\mathcal{C}) \leq s$. By Claim 1, $|\mathcal{C}| = m(s) \leq s+1$. If $|\mathcal{C}| \leq s$, then $o(\mathcal{C}) \leq s$. Assume that $|\mathcal{C}| = s+1$. Since $P = S \cup \{\bigcup_{i=1}^{s+1} V(C_i)\}$, the total number of vertices of P is equal to $n = s + \sum_{i=1}^{s+1} |V(C_i)|$. Consider two cases where (i) s is odd, (ii) s is even. In both cases if all the components in \mathcal{C} are odd, then n is odd; contradicting our assumption that P has an even number of vertices. Thus, \mathcal{C} contains at least one even component, which implies that $o(\mathcal{C}) \leq s$.

Finally, by Claim 2 and Theorem 1, we conclude that 2-GG has a perfect matching. \square

Theorem 6. For every set P of n points, 1-GG has a matching of size at least $\frac{2(n-1)}{5}$.

Proof. Let S be a set of s vertices removed from 1-GG, and let $\mathcal{C} = \{C_1, \dots, C_{m(s)}\}$ be the resulting $m(s)$ components. Actually $\mathcal{C} = 1\text{-GG} - S$ and $\mathcal{P} = \{V(C_1), \dots, V(C_{m(s)})\}$ is a partition of the vertices in $P \setminus S$. Note that $o(\mathcal{C}) \leq m(s)$. Let M^* be a maximum matching in 1-GG. By Theorem 2,

$$|M^*| = \frac{1}{2}(n - \text{def}(1-GG)), \quad (2)$$

where

$$\begin{aligned} \text{def}(1-GG) &= \max_{S \subseteq P} (o(\mathcal{C}) - |S|) \\ &\leq \max_{S \subseteq P} (|\mathcal{C}| - |S|) \\ &= \max_{0 \leq s \leq n} (m(s) - s). \end{aligned} \quad (3)$$

Define $G(\mathcal{P})$, \mathcal{T} , \mathcal{D} , and F as in the proof of Theorem 5. By Theorem 4, $|F| \leq 3 \cdot |S|$. By the same reasoning as in the proof of Theorem 5, each disk in \mathcal{D} has at least two points of S in its interior. Thus, $2 \cdot |\mathcal{D}| \leq |F|$. Therefore, $2(m(s) - 1) \leq |F| \leq 3s$, and hence

$$m(s) \leq \frac{3s}{2} + 1. \quad (4)$$

In addition, $s + m(s) = |S| + |\mathcal{C}| \leq |P| = n$, and hence

$$m(s) \leq n - s. \quad (5)$$

By Inequalities (4) and (5),

$$m(s) \leq \min\left\{\frac{3s}{2} + 1, n - s\right\}. \quad (6)$$

Thus, by (3) and (6)

$$\begin{aligned} \text{def}(1-GG) &\leq \max_{0 \leq s \leq n} (m(s) - s) \\ &\leq \max_{0 \leq s \leq n} \left\{ \min\left\{\frac{3s}{2} + 1, n - s\right\} - s \right\} \\ &= \max_{0 \leq s \leq n} \left\{ \min\left\{\frac{s}{2} + 1, n - 2s\right\} \right\} \\ &= \frac{n + 4}{5}, \end{aligned} \quad (7)$$

where the last equation is achieved by setting $\frac{s}{2} + 1$ equal to $n - 2s$, which implies $s = \frac{2(n-1)}{5}$. Finally by substituting (7) in Equation (2) we have

$$|M^*| \geq \frac{2(n-1)}{5}.$$

□

By similar reasoning as in the proof of Theorem 6 we have the following Theorem.

Theorem 7. *For every set P of n points, 0 - GG has a matching of size at least $\frac{n-1}{4}$.*

The bound in Theorem 7 is tight, as can be seen from the graph in Figure 9, for which the maximum matching has size $\frac{n-1}{4}$. Actually this is a Gabriel graph of maximum degree four which is a tree. The dashed edges do not belong to 0 - GG because any closed disk which has one of these edges as diameter has a point on its boundary. Observe that each edge in any matching is adjacent to one of the vertices of degree four.




Figure 9: A 0- GG of $n = 17$ points with a maximum matching of size $\frac{n-1}{4} = 4$ (bold edges). The dashed edges do not belong to the graph because any of their corresponding closed disks has a point on its boundary.

Note: For a point set P , let $\nu_k(P)$ and $\alpha_k(P)$ respectively denote the size of a maximum matching and a maximum independent set in k - GG . For every edge in the maximum matching, at most one of its endpoints can be in the maximum independent set. Thus,

$$\alpha_k(P) \leq |P| - \nu_k(P).$$

By combining this formula with the results of Theorems 7, 6, 5, respectively, we have $\alpha_0(P) \leq \frac{3n+1}{4}$, $\alpha_1(P) \leq \frac{3n+2}{5}$, and $\alpha_2(P) \leq \lceil \frac{n}{2} \rceil$. The 0- GG graph in Figure 9 has an independent set of size $\frac{3n+1}{4} = 13$, which shows that this bound is tight for 0- GG . On the other hand, 0- GG is planar and every planar graph is 4-colorable; which implies that $\alpha_0(P) \geq \lceil \frac{n}{4} \rceil$. There are some examples of 0- GG in [13] such that $\alpha_0(P) = \lceil \frac{n}{4} \rceil$, which means that this bound is tight as well.

5 Blocking Higher-Order Gabriel Graphs

In this section we consider the problem of blocking higher-order Gabriel graphs. Recall that a point set K blocks k - $GG(P)$ if in k - $GG(P \cup K)$ there is no edge connecting two points in P .

Theorem 8. *For every set P of n points, at least $\lceil \frac{n-1}{3} \rceil$ points are necessary to block 0- $GG(P)$.*

Proof. Let K be a set of m points which blocks 0- $GG(P)$. Let $G(\mathcal{P})$ be the complete graph with vertex set $\mathcal{P} = P$. Let \mathcal{T} be a minimum spanning tree of $G(\mathcal{P})$ and let \mathcal{D} be the set of closed disks corresponding to the edges of \mathcal{T} . Since $G(\mathcal{P})$ has n vertices, \mathcal{T} has $n - 1$ edges. Thus, $|\mathcal{D}| = n - 1$. By Lemma 3 each disk $D[a, b] \in \mathcal{D}$ does not contain any point of $P \setminus \{a, b\}$, thus, $\mathcal{T} \subseteq 0$ - $GG(P)$. To block each edge of \mathcal{T} , corresponding to a disk in \mathcal{D} , at least one point is necessary. By Theorem 4 each point in K can lie in at most three disks of \mathcal{D} . Therefore, $m \geq \lceil \frac{n-1}{3} \rceil$, which implies that at least $\lceil \frac{n-1}{3} \rceil$ points are necessary to block all the edges of \mathcal{T} and hence 0- $GG(P)$. \square

Figure 10(a) shows a 0- GG with $n = 13$ (black) points which is blocked by $\lceil \frac{n-1}{3} \rceil = 4$ (white) points. Note that all the disks, corresponding to the edges of every cycle, intersect at the same point in the plane (where we have placed the white points). As shown in Figure 10(b), the dashed edges do not belong to 0- GG . Thus, the lower bound provided by Theorem 8 is tight. It is easy to generalize the result of Theorem 8 to higher-order Gabriel graphs. Since in a k - GG we need at least $k + 1$ points to block an edge of \mathcal{T} and each point can be inside at most three disks in \mathcal{D} , we have the following corollary:

Corollary 2. *For every set P of n points, at least $\lceil \frac{(k+1)(n-1)}{3} \rceil$ points are necessary to block k - $GG(P)$.*

In [3] the authors showed that every Gabriel graph can be blocked by a set K of $n - 1$ points by putting a point slightly to the right of each point of P , except for the rightmost one. Every




Figure 10: (a) 0- GG graph of $n = 13$ points (in bold edges) which is blocked by $\lceil \frac{n-1}{3} \rceil = 4$ white points, (b) dashed edges do not belong to 0- GG .

disk with diameter determined by two points of P will contain a point of K . Using a similar argument one can block a k - GG by putting $k + 1$ points slightly to the right of each point of P , except for the rightmost one. Thus,

Corollary 3. *For every set P of n points, there exists a set of $(k + 1)(n - 1)$ points that blocks k - $GG(P)$.*

Note that this upper bound is tight, because if the points of P are on a line, the disks representing the minimum spanning tree are disjoint and each disk needs $k + 1$ points to block the corresponding edge.

6 Conclusion

In this paper, we considered the bottleneck and perfect matching admissibility of higher-order Gabriel graphs. We proved that

- 10- GG contains a Euclidean bottleneck matching of P and 8- GG may not have any.
- 0- GG has a matching of size at least $\frac{n-1}{4}$ and this bound is tight.
- 1- GG has a matching of size at least $\frac{2(n-1)}{5}$.
- 2- GG has a perfect matching.
- At least $\lceil \frac{n-1}{3} \rceil$ points are necessary to block 0- GG and this bound is tight.
- $\lceil \frac{(k+1)(n-1)}{3} \rceil$ points are necessary and $(k + 1)(n - 1)$ points are sufficient to block k - GG .

We leave a number of open problems:

- Does 9- GG contain a Euclidean bottleneck matching of P ?
- What is a tight lower bound on the size of a maximum matching in 1- GG ?

References

- [1] M. Abellanas, P. Bose, J. García-López, F. Hurtado, C. M. Nicolás, and P. Ramos. On structural and graph theoretic properties of higher order Delaunay graphs. *Int. J. Comput. Geometry Appl.*, 19(6):595–615, 2009.

- [2] O. Aichholzer, R. F. Monroy, T. Hackl, M. J. van Kreveld, A. Pilz, P. Ramos, and B. Vogtenhuber. Blocking Delaunay triangulations. *Comput. Geom.*, 46(2):154–159, 2013.
- [3] B. Aronov, M. Dulieu, and F. Hurtado. Witness Gabriel graphs. *Comput. Geom.*, 46(7):894–908, 2013.
- [4] J. Babu, A. Biniiaz, A. Maheshwari, and M. Smid. Fixed-orientation equilateral triangle matching of point sets. To appear in *Theoretical Computer Science*.
- [5] C. Berge. Sur le couplage maximum d’un graphe. *C. R. Acad. Sci. Paris*, 247:258–259, 1958.
- [6] A. Biniiaz, A. Maheshwari, and M. Smid. Matching in Gabriel graphs. arXiv:1410.0540 [cs.CG], 2014.
- [7] P. Bose, S. Collette, F. Hurtado, M. Korman, S. Langerman, V. Sacristan, and M. Saumell. Some properties of k -Delaunay and k -Gabriel graphs. *Comput. Geom.*, 46(2):131–139, 2013.
- [8] M.-S. Chang, C. Y. Tang, and R. C. T. Lee. 20-relative neighborhood graphs are Hamiltonian. *Journal of Graph Theory*, 15(5):543–557, 1991.
- [9] M.-S. Chang, C. Y. Tang, and R. C. T. Lee. Solving the Euclidean bottleneck matching problem by k -relative neighborhood graphs. *Algorithmica*, 8(3):177–194, 1992.
- [10] M. B. Dillencourt. Toughness and Delaunay triangulations. *Discrete & Computational Geometry*, 5:575–601, 1990.
- [11] K. R. Gabriel and R. R. Sokal. A new statistical approach to geographic variation analysis. *Systematic Zoology*, 18(3):259–278, 1969.
- [12] T. Kaiser, M. Saumell, and N. V. Cleemput. 10-Gabriel graphs are Hamiltonian. arXiv:1410.0309v3 [cs.CG], 2015.
- [13] D. W. Matula and R. R. Sokal. Properties of Gabriel graphs relevant to geographic variation research and the clustering of points in the plane. *Geographical Analysis*, 12(3):205–222, 1980.
- [14] T.-H. Su and R.-C. Chang. The k -Gabriel graphs and their applications. In *SIGAL International Symposium on Algorithms*, pages 66–75, 1990.
- [15] W. T. Tutte. The factorization of linear graphs. *Journal of the London Mathematical Society*, 22(2):107–111, 1947.